

**SOCIAL NORMS MARKETING:
THE LOGIC AND NECESSARY DATA
TO DEMONSTRATE EFFECTIVENESS**



<http://www.socialnorms.org>

**NATIONAL INSTITUTE FOR SOCIAL NORMS
UNIVERSITY OF VIRGINIA
CHARLOTTESVILLE, VA 22903**

AUTHORS:

**ADRIENNE KELLER, PH.D., RESEARCH DIRECTOR
JENNIFER BAUERLE, PH.D., DIRECTOR
JAMES TURNER, M.D., EXECUTIVE DIRECTOR**

FEBRUARY 2008

Contents

Introduction

The hypothesized causal pathway connecting social norms marketing campaigns to harm reduction

Model of the logic of social norms marketing campaigns and associated evidence

Relationship of model components to questions included in the NCHA

Overview of the methodology of a social norms marketing campaign

Brief tool to assist in evaluation of the process and impact of social norms marketing campaigns

Some helpful references


Introduction

The materials that follow were developed at NSNI as a working tool primarily to assist colleges and universities who are conducting social norms marketing interventions to reduce the adverse consequences associated with high risk alcohol consumption. Since many colleges and universities participate in the National College Health Assessment (NCHA), this document uses questions in the NCHA as examples of available relevant information.

The model and the assessment tool, although both written in terms of social norms marketing campaigns directed towards alcohol consumption, can be adapted for other issues. Please contact us at NSNI if you would like assistance in adapting this material for other social norms marketing interventions.

The hypothesized causal pathway


The social norms model hypothesizes that, against a background of a wide variety of influences (Box 1), a person's perceptions of peers' behaviors and beliefs surrounding alcohol (Box 2) will have a strong influence on both personal drinking patterns (Box 3a) and use of protective strategies (Box 3b). These will, in turn, strongly influence the number, type and severity of adverse consequences associated with hazardous drinking (Box 4).




*Characteristics in bold are measured by variables in the National College Health Assessment (NCHA) survey conducted of the American College Health Association (<http://www.acha-ncha.org/>)

A *social norms marketing intervention* aims to correct the misperception of peers' behavior in order to influence personal choices. The messages typically focus on the reality of peers' drinking but may also include the use of protective behaviors by peers. For social norms marketing interventions, the causal pathway can be used to create a logic model that relates the intervention to goals, outcomes and impact.

LOGIC OF SOCIAL NORMS INTERVENTIONS



EVIDENCE NEEDED



*If norms for protective behaviors were included in the marketing campaign

**Baseline assessment needed before intervention & follow-up assessment after intervention

Necessary Evidence to Demonstrate Effectiveness: What's Included in the NCHA

Sample Description (*NCHA questions 45 through 58*)

Evidence related to underlying problems

- Perception of norms related to drinking (*NCHA questions 10d, 15, 19b*)
- Drinking patterns (*NCHA questions 9d, 12, 13, 14, 16, and calculated eBAC*)

Evidence related to intervention strategy (*not included in NCHA*)

- Fidelity: Marketing materials conform to social norms principles
 - Materials address correcting a significant misperception related to peers' behaviors
 - Conforms to PIE principles: Positive, Inclusive, Empowering
- Reach: Marketing reaches intended audience
 - "Exposure": evidence that a majority of the intended audience was exposed to the campaign
 - "Saturation": evidence that information was seen a sufficient number of times by intended audience
 - Suggested question to measure exposure: *In the past [time frame], how many times have you seen [or heard] the [campaign logo/brand or message or specific campaign material]?*

Evidence related to intervention goal

- There is a significant improvement from pre-intervention to post-intervention in measure(s) of normative perceptions:
 - Descriptive (*NCHA questions 10d, 15, 19b*)
 - Injunctive (*not included in NCHA*)
 - Example of injunctive norm: Most AnywhereU students approve of drinking to get drunk.

Evidence related to anticipated outcomes

- There is a significant improvement from pre-intervention to post-intervention in measure(s) of protective behaviors (*may not happen if campaign did not focus on protective behaviors*):
 - Personal protective behaviors (*NCHA question 17a through 17j*)
 - Prevalence
 - Frequency of use
 - Protective behaviors for others (*not included in NCHA*)
 - Prevalence
 - Frequency of use
- There is a significant improvement from pre-intervention to post-intervention in measure(s) of patterns of drinking; for example:
 - Number of drinks consumed last time partied or socialized (*NCHA question 13*)
 - Hours over which the drinks were consumed (*NCHA question 12*)
 - eBAC: Above two are used, with sex (*NCHA question 46*), weight (*NCHA question 48*) and gender constant, in equation to compute estimated blood alcohol content (eBAC)
 - Number of days in previous month drank alcohol (*NCHA question 9d*)
 - Times in last two weeks drank as much or more than last time partied/socialized (*NCHA question 14*)
 - Times drank more than 5 drinks (*NCHA question 16*)

Evidence related to desired impact

- There is a significant improvement from pre-intervention to post-intervention in measure(s) of adverse consequences associated with drinking (*NCHA question 18a through 18g: 7 adverse consequences*)

Optional additional helpful evidence

- Level of drinking and negative consequences did NOT increase from pre-intervention to post-intervention among low risk drinkers
- Perceptions, drinking and negative consequences all changed in the desired direction from pre-intervention to post-intervention among high risk drinkers

Overview of Methodology

- Conduct a (stratified) random sample survey of intended population
 - Ensure sufficient sample size for sufficient power to examine subgroups (e.g., by sex or class level) taking into account likely response rate
 - Clean the data (eliminate incomplete essential data, artifactual data)
 - Balance sample demographic characteristics to match population characteristics, for example:
 - Sex
 - Academic class level (in school populations)
 - Race/ethnicity
- Establish the baseline rate of behaviors and attitudes of interest:
 - Quantity/frequency of drinking: a variety of possible variables including drinks per week, drinks per weekend, drinks last time partied/socialized, eBAC
 - Perception of drinking of others: perception questions *must* correspond exactly to quantity/frequency questions
 - Use of protective behaviors (may be omitted)
 - Adverse consequences associated with drinking
- Test if there is a *mis*perception of the true drinking norm: are the perceived quantity/frequency measures significantly greater than the actual quantity/frequency measures?
- Test if this misperception is associated with hazardous drinking
 - Those who drink more are more likely to perceive the drinking norm to be greater than the true drinking norm, [and close to or greater than what they themselves drink, per Berkowitz]
 - Establish the percent of the population that have normative misperceptions *and* hazardous drinking
- Design a social norms marketing campaign
 - Use representatives from intended population to design message and marketing strategies
 - Use a variety of messages and/or message delivery strategies
- Test if the marketing campaign *reached* the intended audience
 - Exposure: What percentage of the population saw the messages?
 - Saturation: How frequently did members of the intended population see the message?
- Conduct a follow-up (stratified) random sample survey of the intended population:
 - Same questions about drinking behaviors and attitudes, protective behaviors and adverse consequences as in the first step
 - Closely following conclusion of campaign
 - Clean and balance sample data as in the first step
- Test if observed changes conformed to theory:
 - Is there a decrease in perception of drinking norms?
 - Is there a decrease in actual drinking?
 - Is there an increase in protective behaviors (if included as focus of campaign)?
 - Is there a decrease in adverse consequences?

Evaluation Tool Assessing the Strength of Necessary Evidence and Inferences:

This tool is designed to help you assess the strengths of the six essential components of the model. Remember that the logic connecting your intervention to observed changes is only as strong as your *weakest* component. Ideally, each area should score at least 8; if any area scores less than 6 that is cause for concern. The overall score should be at least 49; overall scores less than 38 indicate that your evidence is probably not strong enough to demonstrate effectiveness convincingly.

1. The data are valid and reliable:	Completely	Mostly	Somewhat	Not at all or information not available
a. The sampling strategy was unbiased.	3	2	1	0
b. The resultant sample matched the population in key descriptive variables	3	2	1	0
c. All variables of interest were assessed using standardized measures with demonstrated validity and reliability in the population of interest	3	2	1	0
Total Score for Data Quality:				
2. Hazardous drinking and misperceptions of norms are underlying problems :	Agreement among all measures	Agreement among most measures	Agreement among some but not most measures	Not at all or information not available
a. Hazardous drinking patterns are prevalent:	3	2	1	0
b. Misperceptions of drinking norms are prevalent:	3	2	1	0
c. Positive correlation between hazardous drinking patterns and misperceptions:	3	2	1	0
Total Score for Evidence of Underlying Problems:				
3. The intervention was well implemented:	Completely	Mostly	Somewhat	Not at all or information not available
a. The social norms campaign conformed to social norms marketing principles	3	2	1	0
b. The intended audience was exposed to the campaign:	3	2	1	0
c. Most members of the intended audience had at least 5-10 exposures	3	2	1	0
Total Score for Evidence of Well-Implemented Intervention:				

4. Intervention Goal *: Following the campaign, accuracy of perceptions increased:	Change is statistically significant and clearly meaningful	Change is small but statistically significant	Change occurred but is not statistically significant	No evidence, no change, or change in wrong direction
a. The perceptions of norms that were a focus of the campaign were more accurate:	3	2	1	0
b. The perception of norms that were <i>not</i> a focus of the campaign did <i>not</i> improve in accuracy	3	2	1	0
c. High-risk populations (e.g., binge drinkers) increased accuracy of perceptions	3	2	1	0
Total Score for Evidence that Intervention Goal was Achieved:				<input type="text"/>
5. Anticipated Outcomes *: Following the campaign, decreased <i>risk</i> for harm was observed:	Change is statistically significant and clearly meaningful	Change is small but statistically significant	Change occurred but is not statistically significant	No evidence, no change, or change in wrong direction
a. Hazardous drinking behaviors decreased overall	3	2	1	0
b. eBAC decreased overall	3	2	1	0
c. High-risk populations (e.g., binge drinkers) experienced decreased risk for harm	3	2	1	0
Total Score for Evidence that Intervention Outcomes were Achieved:				<input type="text"/>
6. Desired Impact *: Following the campaign, significant harm reduction was observed:	Change is statistically significant and clearly meaningful	Change is small but statistically significant	Change occurred but is not statistically significant	No evidence, no change, or change in wrong direction
a. There is a decrease in adverse consequences associated with drinking:	3	2	1	0
b. There is an increase in percent experiencing no adverse consequences	3	2	1	0
c. High-risk populations (e.g., binge drinkers) experienced fewer adverse consequences	3	2	1	0
Total Score for Evidence that the Desired Impact was Achieved:				<input type="text"/>
Overall Total Score for Necessary Evidence & Inferences:				<input type="text"/>

*For multi-year projects, the measures of perceptions, behaviors and consequences following each year of intervention become the baseline measures for the next year.

Some Key References

- Bauerle, J (2003). The University of Virginia's Social Norms Marketing Campaign. *The Report on Social Norms: Working Paper #11*. Little Falls, NJ: PaperClip Communications.
- Bauerle J, Burwell C & Turner JC (2002). Social Norms Marketing at the University of Virginia. Presented at the Annual Meeting of the American College Health Association, May 23, Washington, DC.
- Berkowitz AD (2002). Responding to the Critics: Answers to Common Questions and Concerns About the Social Norms Approach. *The Report on Social Norms: Working Paper #7*. Little Falls, NJ: PaperClip Communications.
- Borsari B & Carey KB (2003). Descriptive and injunctive norms in college drinking: a meta-analytic integration. *Journal of Studies on Alcohol*, 64(3), 331-41.
- Bruce S, Hawkins P, Sharp M & Keller A (2006). Experience-Based Suggestions for Achieving a High Survey Response Rate. *Report on Social Norms*, <http://www.mostofus.org/newsletter/index.php?newsletterID=5>.
- Bruce S & Keller A (2007). Applying Social Norms Theory within Affiliation Groups: Promising Interventions for High-Risk Drinking, *NASPA JI* 44(1), 101-107.
- Haines MP, Perkins HW, Rice RM & Barker G (2005). A Guide to Marketing Social Norms for Health Promotion in Schools and Communities, <http://www.socialnormsresources.org/pdf/Guidebook.pdf>.
- Mattern JL & Neighbors C (2004). Social norms campaigns: examining the relationship between changes in perceived norms and changes in drinking levels. *Journal of Studies on Alcohol*, 65(4), 489-93.
- Perkins HW (2002). Social norms and the prevention of alcohol misuse in collegiate contexts. *Journal of Studies on Alcohol, Supplement* (14), 164-72.
- Perkins HW & Berkowitz AD (1986). *Perceiving the Community Norms of Alcohol Use Among Students: Some Research Implications for Campus Alcohol Education Programming*. *International Journal of the Addictions*, 21(9/10), 961-976.
- Turner JC, Bauerle J & Shu J (2004). Estimated blood alcohol concentration correlation with self-reported negative consequences among college students using alcohol. *Journal of Studies on Alcohol*, 741-749.